



Infographic



Journals
BAHIANA
SCHOOL OF MEDICINE AND PUBLIC HEALTH



Effects and parameters of taVNS in individuals with atrial fibrillation: an infographic

Maria Williane de Sousa Ribeiro¹

Nicole Beatriz Nascimento Miranda²

Fernanda Leite Dias Dantas Estevam³

Larissa Conceição Dias Lopes⁴

Tiago da Silva Lopes⁵

¹⁻³Colégio Adventista da Bahia (Cachoeira). Bahia, Brazil.

⁴Universidade Federal da Bahia (Salvador). Bahia, Brazil.

⁵Corresponding author. Universidade Federal do ABC (São Bernardo do Campo). São Paulo, Brazil. tslopes.physio@gmail.com

ABSTRACT | BACKGROUND: Transcutaneous auricular vagus nerve stimulation (taVNS) is a non-invasive neuromodulatory technique used to modulate autonomic excitability through the electrical stimulus of the vagus nerve. Studies have shown that taVNS can be used as a therapeutic strategy for arterial fibrillation (AF) management. This infographic aims to highlight the evidence about the effect and parameters of taVNS in individuals with AF. **METHOD:** We conducted a review using the PubMed mesh combination (Transcutaneous Electric Nerve Stimulation)AND(Nerve Stimulation, Vagal)OR(Nerve Stimulation, Vagus)OR(Nerve Stimulation, Vagal)AND(Atrial Fibrillation) that included six taVNS clinical trials with human adults over the age of 18. **RESULTS:** taVNS can promote an increase in the PQ interval on the electrocardiogram by increasing the parasympathetic tone. Moreover, the taVNS can prevent AF progression through antiarrhythmic effects, such as antiadrenergic mechanisms of neural remodeling, and a decrease of pro-inflammatory cytokines and AF of burden. **DISCUSSION:** The acute effect of taVNS can increase the P - wave alternans (PWA), which is believed to be related to AF. However, the chronic effect of taVNS for six months promotes a decrease in PWA and AF burden. In addition, the taVNS can alter the heart variability ratio parameters through the increase of the HF/LF relation in individuals with AF, which can be a signal of sympathovagal rebalance. However, the taVNS can improve the autonomic balance through the decrease in HF/LH relation in healthy individuals. **CONCLUSION:** The taVNS can be a useful therapeutic tool in the management of individuals with AF due to its antiarrhythmic effects.

KEYWORDS: taVNS. NIBS. Neuromodulation. Persistent Atrial Fibrillation.

Submitted 10/05/2022, Accepted 03/30/2023, Published 05/12/2023

Brain Imaging Stimul., Salvador, 2023;2:e4871

<http://dx.doi.org/10.17267/2965-3738bis.2023.e4871>

ISSN: 2965-3738


Assigned editor: Ana Paula Fontana, Katia Nunes Sá

How to cite this article: Ribeiro MWS, Miranda NBN, Estevam FLDD, Lopes LCD, Lopes TS. Effects and parameters of taVNS in individuals with atrial fibrillation: an infographic. Brain Imaging Stimul.




2023;2:e4871. <http://dx.doi.org/10.17267/2965-3738bis.2023.e4871>




Effects and Parameters of taVNS in Atrial Fibrillation



(1)



(1,5,6)



Infographic Legend

taVNS can be applied on the ear tragus or cymba concha using two electrodes, a cathode, and an anode. Some studies^{5,7} have been using electrodes in the ear lobule as the control condition due to the lack of vagus innervation in this region.⁴ The antiarrhythmogenic effects of taVNS include the antiadrenergic mechanisms of neural remodeling and a decrease of pro-inflammatory cytokines and AF of burden.⁸⁻¹⁰ This infographic layout was built using Biorender.com in agreement with G.R.A.P.H.I.C guidelines.¹¹

Authors' contributions

All authors were involved in all aspects of infographic conception. Ribeiro MWS drafted the layout infographic and wrote the abstract text, while Lopes TS, Lopes LCD, Estevam FLDD and Miranda NBN critically reviewed it.

Conflicts of interest

No financial, legal, or political conflicts involving third parties (government, companies, and private foundations, etc.) were declared for any aspect of the submitted work (including, but not limited to grants and funding, participation in an advisory board, study design, preparation manuscript, statistical analysis, etc.).

References

1. Kharbanda RK, Does WFB, Staveten LN, Taverne YJHJ, Bogers AJJC, Groot NMS. Vagus Nerve Stimulation and Atrial Fibrillation: Revealing the Paradox. *Neuromodulation*. 2022;25(3):356-365. <https://doi.org/10.1016/j.neurom.2022.01.008>
2. Badran BW, Dowdle LT, Mithoefer OJ, LaBate NT, Coatsworth J, Brown JC, et al. Neurophysiologic Effects of Transcutaneous Auricular Vagus Nerve Stimulation (taVNS) via Electrical Stimulation of the Tragus: A Concurrent taVNS/fMRI Study and Review. *Focus*. 2022;20(1):80-9. <https://doi.org/10.1176/appi.focus.20110>
3. Clancy JA, Mary DA, Witte KK, Greenwood JP, Deuchars SA, Deuchars J. Non-invasive vagus nerve stimulation in healthy humans reduces sympathetic nerve activity. *Brain Stimul*. 2014;7(6):871-7. <https://doi.org/10.1016/j.brs.2014.07.031>
4. Peuker ET, Filler TJ. The nerve supply of the human auricle. *Clin Anat*. 2002;15(1):35-7. <https://doi.org/10.1002/ca.1089>
5. Stavrakis S, Humphrey MB, Scherlag BJ, Hu Y, Jackman WM, Nakagawa H, et al. Low-level transcutaneous electrical vagus nerve stimulation suppresses atrial fibrillation. *J Am Coll Cardiol*. 2015;65(9):867-75. <https://doi.org/10.1016/j.jacc.2014.12.026>
6. Stavrakis S, Stoner JA, Humphrey MB, Morris L, Filiberti A, Reynolds JC, et al. TREAT AF (Transcutaneous Electrical Vagus Nerve Stimulation to Suppress Atrial Fibrillation): A Randomized Clinical Trial. *JACC Clin Electrophysiol*. 2020;6(3):282-91. <https://doi.org/10.1016/j.jacep.2019.11.008>
7. Chen M, Wang S, Li X, Yu L, Yang H, Liu Q, et al. Non-invasive Autonomic Neuromodulation Is Opening New Landscapes for Cardiovascular Diseases. *Front Physiol*. 2020;11:550578. <https://doi.org/10.3389/fphys.2020.550578>
8. Dagleish AS, Kania AM, Stauss HM, Jelen AZ. Occipitotlantal decompression and noninvasive vagus nerve stimulation slow conduction velocity through the atrioventricular node in healthy participants. *J Osteopath Med*. 2021;121(4):349-59. <https://doi.org/10.1515/jom-2020-0213>
9. Andreas M, Arzl P, Mitterbauer A, Ballarini NM, Kainz FM, Kocher A, et al. Electrical Stimulation of the Greater Auricular Nerve to Reduce Postoperative Atrial Fibrillation. *Circ Arrhythm Electrophysiol*. 2019;12(10):e007711. <https://doi.org/10.1161/CIRCEP.119.007711>
10. Kulkarni K, Singh JP, Parks KA, Katritsis DG, Stavrakis S, Armoundas AA. Low-Level Tragus Stimulation Modulates Atrial Alternans and Fibrillation Burden in Patients With Paroxysmal Atrial Fibrillation. *J Am Heart Assoc*. 2021;10(12):e020865. <https://doi.org/10.1161/JAHA.120.020865>
11. Stones C, Gent M. *The 7 G.R.A.P.I.C Principles of Public Health Infographic Design*. Leeds: University of Leeds; 2015.